

ISSN 2224-5286

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН

NEWS

OF THE ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN

**ХИМИЯ ЖӘНЕ ТЕХНОЛОГИЯ
СЕРИЯСЫ**



**СЕРИЯ
ХИМИИ И ТЕХНОЛОГИИ**



**SERIES
CHEMISTRY AND TECHNOLOGY**

6 (414)

**ҚАРАША – ЖЕЛТОҚСАН 2015 ж.
НОЯБРЬ – ДЕКАБРЬ 2015 г.
NOVEMBER – DECEMBER 2015**

1947 ЖЫЛДЫҢ ҚАҢТАР АЙЫНАН ШЫҒА БАСТАҒАН
ИЗДАЕТСЯ С ЯНВАРЯ 1947 ГОДА
PUBLISHED SINCE JANUARY 1947

ЖЫЛЫНА 6 РЕТ ШЫҒАДЫ
ВЫХОДИТ 6 РАЗ В ГОД
PUBLISHED 6 TIMES A YEAR

АЛМАТЫ, ҚР ҰҒА
АЛМАТЫ, НАН РК
ALMATY, NAS RK

Б а с р е д а к т о р

ҚР ҰҒА академигі
М. Ж. Жұрынов

Р е д а к ц и я а л қ а с ы:

хим. ғ. докторы, проф., ҚР ҰҒА академигі **Әдекенов С.М.**; хим. ғ. докторы, проф., ҚР ҰҒА академигі **Ғазалиев А.М.**; хим. ғ. докторы, проф., ҚР ҰҒА академигі **Ерғожин Е.Е.** (бас редактордың орынбасары); хим. ғ. докторы, проф., ҚР ҰҒА академигі **Пірәлиев К.Д.**; хим. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Баешов А.Б.**; хим. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Бүркітбаев М.М.**; хим. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Жүсіпбеков У.Ж.**; хим. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Итжанова Х.И.**; хим. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Молдахметов М.З.**, техн. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Мырхалықов Ж.У.**; мед. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Рахымов К.Д.**; хим. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Сатаев М.И.**; хим. ғ. докторы, проф., ҚР ҰҒА корр. мүшесі **Тәшімов Л.Т.**; хим. ғ. докторы, проф. **Мансұров З.А.**; техн. ғ. докторы, проф. **Наурызбаев М.К.**

Р е д а к ц и я к е ң е с і:

Беларусь Республикасының ҰҒА академигі **Агабеков В.Е.** (Беларусь); Украинаның ҰҒА академигі **Волков С.В.** (Украина); Қырғыз Республикасының ҰҒА академигі **Жоробекова Ш.Ж.** (Қырғызстан); Армения Республикасының ҰҒА академигі **Мангашян А.А.** (Армения); Молдова Республикасының ҰҒА академигі **Туртэ К.** (Молдова); Әзірбайжан ҰҒА академигі **Фарзалиев В.** (Әзірбайжан); Тәжікстан Республикасының ҰҒА академигі **Халиков Д.Х.** (Тәжікстан); хим. ғ. докторы, проф. **Нараев В.Н.** (Ресей Федерациясы); философия ғ. докторы, профессор **Полина Прокопович** (Ұлыбритания); хим. ғ. докторы, профессор **Марек Сикорски** (Польша)

Главный редактор

академик НАН РК

М. Ж. Журинов

Редакционная коллегия:

доктор хим. наук, проф., академик НАН РК **С.М. Адекенов**; доктор хим. наук, проф., академик НАН РК **А.М. Газалиев**; доктор хим. наук, проф., академик НАН РК **Е.Е. Ергожин** (заместитель главного редактора); доктор хим. наук, проф., академик НАН РК **К.Д. Пралиев**; доктор хим. наук, проф., чл.-корр. НАН РК **А.Б. Башов**; доктор хим. наук, проф., чл.-корр. НАН РК **М.М. Буркитбаев**; доктор хим. наук, проф., чл.-корр. НАН РК **У.Ж. Джусипбеков**; доктор хим. наук, проф., чл.-корр. НАН РК **Х.И. Итжанова**; доктор хим. наук, проф., чл.-корр. НАН РК **М.З. Мулдахметов**; доктор техн. наук, проф., чл.-корр. НАН РК **Ж.У. Мырхалыков**; доктор мед. наук, проф., чл.-корр. НАН РК **К.Д. Рахимов**; доктор хим. наук, проф., чл.-корр. НАН РК **М.И. Сатаев**; доктор хим. наук, проф., чл.-корр. НАН РК **Л.Т. Ташимов**; доктор хим. наук, проф. **З.А. Мансуров**; доктор техн. наук, проф. **М.К. Наурызбаев**

Редакционный совет:

академик НАН Республики Беларусь **В.Е. Агабеков** (Беларусь); академик НАН Украины **С.В. Волков** (Украина); академик НАН Кыргызской Республики **Ш.Ж. Жоробекова** (Кыргызстан); академик НАН Республики Армения **А.А. Манташян** (Армения); академик НАН Республики Молдова **К. Туртэ** (Молдова); академик НАН Азербайджанской Республики **В. Фарзалиев** (Азербайджан); академик НАН Республики Таджикистан **Д.Х. Халиков** (Таджикистан); доктор хим. наук, проф. **В.Н. Нараев** (Россия); доктор философии, профессор **Полина Прокопович** (Великобритания); доктор хим. наук, профессор **Марек Сикорски** (Польша)

«Известия НАН РК. Серия химии и технологии». ISSN 2224-5286

Собственник: Республиканское общественное объединение «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №10893-Ж, выданное 30.04.2010 г.

Периодичность: 6 раз в год

Тираж: 300 экземпляров

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел. 272-13-19, 272-13-18,
<http://nauka-nanrk.kz/chemistry-technology.kz>

© Национальная академия наук Республики Казахстан, 2015

Адрес редакции: 050100, г. Алматы, ул. Кунаева, 142,
Институт органического катализа и электрохимии им. Д. В. Сокольского,
каб. 310, тел. 291-62-80, факс 291-57-22, e-mail:orgcat@nursat.kz

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75

Editor in chief

M. Zh. Zhurinov,
academician of NAS RK

Editorial board:

S.M. Adekenov, dr. chem. sc., prof., academician of NAS RK; **A.M. Gazaliev**, dr. chem. sc., prof., academician of NAS RK; **Ye.Ye. Yergozhin**, dr. chem. sc., prof., academician of NAS RK (deputy editor); **K.D. Praliyev**, dr. chem. sc., prof., academician of NAS RK; **A.B. Bayeshov**, dr. chem. sc., prof., corr. member of NAS RK; **M.M. Burkhitbayev**, dr. chem. sc., prof., corr. member of NAS RK; **U.Zh. Zhusipbekov**, dr. chem. sc., prof., corr. member of NAS RK; **Kh.I. Itzhanova**, dr. chem. sc., prof., corr. member of NAS RK; **M.Z. Muldakhmetov**, dr. eng. sc., prof., corr. member of NAS RK; **Zh.U. Myrkhalykov**, dr. eng. sc., prof., corr. member of NAS RK; **K.D. Rakhimov**, dr. med. sc., prof., corr. member of NAS RK; **M.I. Satayev**, dr. chem. sc., prof., corr. member of NAS RK; **L.T. Tashimov**, dr. chem. sc., prof., corr. member of NAS RK; **Z.A. Mansurov**, dr. chem. sc., prof.; **M.K. Nauryzbayev**, dr. eng. sc., prof.

Editorial staff:

V.Ye. Agabekov, NAS Belarus academician (Belarus); **S.V. Volkov**, NAS Ukraine academician (Ukraine); **Sh.Zh. Zhorobekov**, NAS Kyrgyzstan academician (Kyrgyzstan); **A.A. Mantashyan**, NAS Armenia academician (Armenia); **K. Turte**, NAS Moldova academician (Moldova); **V. Farzaliyev**, NAS Azerbaijan academician (Azerbaijan); **D.Kh. Khalikov**, NAS Tajikistan academician (Tajikistan); **V.N. Narayev**, dr. chem. sc., prof. (Russia); **Pauline Prokopovich**, dr. phylos., prof. (UK); **Marek Sikorski**, dr. chem. sc., prof. (Poland)

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of chemistry and technology.
ISSN 2224-5286

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of Information and Archives of the Ministry of Culture and Information of the Republic of Kazakhstan N 10893-Ж, issued 30.04.2010

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,
<http://nauka-nanrk.kz/chemistry-technology.kz>

© National Academy of Sciences of the Republic of Kazakhstan, 2015

Editorial address: Institute of Organic Catalysis and Electrochemistry named after D. V. Sokolsky
142, Kunayev str., of. 310, Almaty, 050100, tel. 291-62-80, fax 291-57-22,
e-mail: orgcat@nursat.kz

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

SERIES CHEMISTRY AND TECHNOLOGY

ISSN 2224-5286

Volume 6, Number 414 (2015), 109 – 113

UDK 547.792.3

**ALKYLATION AND ACYLATION REACTIONS
OF 3-(MORPHOLINOMETHYL)-4-PHENYL-1H-1,2,4-
TRIAZOLE-5(4H)-THIONE****M. A. Dyussebaeva, J. Jenis, L. B. Zhaimuhambetova, Sh. S. Akhmedova**Al-Farabi Kazakh National University, Almaty, Kazakhstan.
E-mail: zhaimuhambetova@gmail.com**Key words:** morpholine, thiosemicarbazide, triazole, cyclization, S-alkylation, N-acylation.

Abstract. Derivatives of 1,2,4-triazole-5-thiones are heterocyclic compounds with thiol-thione grouping. This assumes tautomeric transformations of 1,2,4-triazoline-5-thione to 1,2,4-triazole-5-thiol. 3-(Morpholinomethyl)-4-phenyl-1H-1,2,4-triazole-5(4H)-thione (II) exists in two tautomeric forms (IIa and IIb). This conducts the reaction of (II) by two reaction centers (by NH or SH groups). The presence of substituents in the 1,2,4-triazole ring promotes the reaction of substitution in the core of 1,2,4-triazole. During the alkylation or acylation of substituted 1,2,4-triazol-5(4H)-thione (II) the acceding group may be attached to either nitrogen atom or a sulfur atom. Thus alkylation with ethyl bromoacetate, methyl iodide and/or butyl bromide in alkaline media gave the corresponding S-alkylated compounds (III), (IV) and (V) respectively. While acylation of triazole (II) with benzoyl chloride in the presence of triethylamine yielded the N-acylated derivative (VI). The structures of the synthesized compounds were confirmed by elemental analyses, IR and H^1 NMR spectra.

Introduction. Search and development of highly biologically active compounds and other practically useful substances and materials is one of the priorities of organic chemistry. In this regard, a promising is the use of a systematic approach, including a targeted synthesis of potentially biologically active substances, the determination of its structure, chemical modification and bioscreening. A decisive role in the chain plays electronic and spatial structure of the synthesized compounds, which determine its reactivity, tendency to tautomeric transformations and biological activity [1].

Among the variety of classes of organic compounds that exhibit a wide range of biological activity leading place saturated heterocycles. Among the heterocyclic derivatives found many hundreds of highly drugs, pesticides, plant growth regulators, and many other substances and materials with application properties. The combination of hydrocarbon fragments and single hetero cyclic or polycyclic system gives those compounds of a number of properties that reflect the particular carbocyclic moiety and a heteroatom functions. Hetero significantly affect the reactivity of neighboring atoms can be included in the π -electron system are structural elements of σ -skeleton. In constructing heterocyclic system often use nitrogen, oxygen, sulfur, phosphorus [2].

Currently, there is an increased interest in the nitrogen-containing heterocyclic compounds having piperidine ring and heterocyclic systems combining different loop nature heteroatoms (morpholine, triazole, thiadiazole, triazolone, oxadiazole, etc.) or different carbo- and heterocycles. This attention is due to the fact that among them found effective painkillers, antispasmodics, antidepressants, anticholinergics, blockers and other biologically active compounds. As a result of research in a number of heterocyclic derivatives, a whole cluster of painkillers, having as a framework the piperidine ring (lidol, morferidin, prodin, promedol, prosidol, kazkain, Rihlokain). Among structurally related compounds found drugs with anti-inflammatory action (flazolone) psychotropic properties (haloperidol, Benperidol). A


compound fused to various cycles, continue the structural number of physiologically active substances - teforin anticholinergic medication, analgesic morphine agonists (oksilarfon) and antagonists (nalorphine, naloxone) [3-6].

In recent years, we had the idea of the piperidine ring as an important component of the fragment of immunomodulation drugs. This, in particular, in the example of the preparation Rihlokain which test data in oncological practice exerts a synergistic effect with respect to cytostatics and acts as an immunomodulator and antitumor drugs prolongator [7].

Among the derivatives of 1,2,4-triazole compounds are also found with antitumor properties, antifungal and sedatives, moreover, a number of compounds known as tranquilizers and substances with analgesic and anti-hypoxic action [1, 8].

Combining piperidine, morpholine fragments with other heterocycles (triazole, oxadiazole) may lead to new reaction centers and changing dietary properties. In the course of our research efforts towards the preparation of new compounds containing the morpholine ring, since several biological activities have reported in morpholine derivatives, have been described here on the synthesis and reactions of new compounds containing the 1, 2, 4-triazole ring as well as morpholine ring.


The starting compound, 3-(morpholinomethyl)-4-phenyl-1*H*-1,2,4-triazole-5(4*H*)-thione (II), was synthesized by the reaction of (morpholil-1-yl)-acetic acid thiosemicarbazide (I) with potassium hydroxide solution [9].



According to IR spectroscopic data of the original compound (II) which has triazolone-5-thione spectre, the observation of $\text{C}=\text{S}$ stretching band of 1240 cm^{-1} and the absence of an absorption about in $2600\text{-}2550\text{ cm}^{-1}$ region cited for SH group have proved that this compound was in the thionic form (IIb) in the solid state. The IR spectrum of triazole (II) showed attributable bands for NH group at 3100 cm^{-1} , for $\text{C}=\text{N}$ group at 1650 cm^{-1} and for aromatic group at $1455\text{-}1500\text{ cm}^{-1}$.

It is well known that alkylation of 1,2,4-triazole-5-thione derivatives with alkyl/arylalkyl halides or alkyl haloester gave the corresponding thioether derivatives. Therefore alkylation of 3-(morpholinomethyl)-4-phenyl-1*H*-1,2,4-triazole-5(4*H*)-thione (II) with ethyl bromo acetate in the presence of potassium carbonate gave ethyl 2-(5-(morpholinomethyl)-4-phenyl-4*H*-1,2,4-triazol-3-ylthio)acetate (III).

The IR spectrum of compound (III) showed attributable bands for $\text{C}=\text{O}$ ester group at 1730 cm^{-1} and for $\text{C}=\text{N}$ group at 1630 cm^{-1} . In the IR spectrum of compound (II), the absorption band of $\text{C}=\text{S}$ group was absent which elucidate that the alkylation reaction was occurred at the sulfur atom.



The ^1H NMR spectrum of compound (III) showed signal at δ 4.0 ppm characteristic for S-CH₂. The ethyl protons resonate at δ 4.1 and 1.3 ppm characteristic for CH₂ protons (quarter).

The absence of signals in the region 4.5 -5.5 ppm elucidate that the alkylation reaction was occurred at the sulfur atom.

Similarly alkylation of (II) with methyl iodide and / or butyl bromide in the presence of triethylamine gave 4-((5-(alkylthio)-4-phenyl-4H-1.2.4-triazole-3-yl)methyl)morpholine (IV) and (V).

Table 1 – The H-NMR spectra of some synthesized compounds in CDCl₃

| Compound (II) | | Compound (III) | | Compound (VI) | |
|-----------------------|----------------|-----------------------|----------------|-----------------------|----------------|
| Group | δ value | Group | δ value | Group | δ value |
| 1H, NH | 11.25 | 5H, Ar | 7.1-7.5 | 10H, Ar-H | 7.3-7.6 |
| 8H, Ar-H | 7.4-7.5 | 2H, N-CH ₂ | 4.09 | 2H, N-CH ₂ | 4.3 |
| 2H, N-CH ₂ | 3.3 | 2H, S-CH ₂ | 4.04 | 4H, 2-CH ₂ | 2.1 |
| 4H, 2-CH ₂ | 2.3 | 4H, 2-CH ₂ | 2.2 | 6H, 3-CH ₂ | 1.7 |
| 6H, 3-CH ₂ | 1.4 | 6H, 3-CH ₂ | 1.5 | | |

The IR spectra of compounds (IV) and (VI) showed attributable bands for C=N groups at 1605-1615 cm⁻¹ and for phenyl groups at 1520-1540 cm⁻¹.

In the IR spectra for the alkylated compounds the absorption bands characteristic for C=S group were absent, this means that the alkylation reaction occurred at the sulfur atom and not at the nitrogen atom.

Acylation reaction of (II) with benzoyl chloride in dry benzene in the presence of triethylamine produced the N-acyl derivative (VI).

The IR spectrum of (3-(morpholinomethyl)-4-phenyl-5-thio-4,5-dihydro-1H-1,2,4-thiazol-1-yl)(phenyl)methanone (VI) showed attributable bands for C=O acyl group at 1730 cm⁻¹, for C=N group at 1620 cm⁻¹, for phenyl group at 1575-1600 cm⁻¹ and for C=S group at 1235 cm⁻¹.

Table 2 – Physical data of the new compounds

| Comp. № | m.p. °C | Formula (M. wt) | Calculated/found | | | | |
|---------|---------|--|------------------|-----------|-------------|-------|-------|
| | | | C | H | N | O | S |
| II | 183-184 | C ₁₃ H ₁₆ N ₄ OS (276.10) | 56.50/56.25 | 5.84/5.66 | 20.27/20.02 | 5.79 | 11.60 |
| III | 254-255 | C ₁₇ H ₂₂ N ₄ O ₃ S (362.14) | 56.33/56.01 | 6.12/5.96 | 15.46/15.76 | 13.24 | 8.85 |
| IV | 109-110 | C ₁₄ H ₁₈ N ₄ OS (290.12) | 57.91/58.04 | 6.25/6.42 | 19.29/19.04 | 5.51 | 11.04 |
| V | 238-240 | C ₁₇ H ₂₄ N ₄ OS (330.43) | 61.41/60.96 | 7.28/7.01 | 16.85/16.55 | 4.81 | 9.64 |
| VI | 240-241 | C ₂₀ H ₂₀ N ₄ O ₂ S (381.43) | 63.14/63.56 | 5.30/5.04 | 14.73/14.21 | 8.41 | 8.43 |

Experimental

Synthesis ethyl 2-(5-(morpholinomethyl)-4-phenyl-4*H*-1,2,4-triazol-3-ylthio)acetate (III). To a solution of triazole (II) (0.01 mole) and anhydrous K₂CO₃ (0.01 mole) in 30 ml acetone was added ethyl bromo acetate (0.011 mole). The reaction mixture was heated at 55-60 °C with stirring for 6 hrs., after filtration the solvent was removed under vacuum and the solid product was filtered off and crystallized from ethanol to give white crystals of ethyl 2-(5-(morpholinomethyl)-4-phenyl-4*H*-1,2,4-triazol-3-ylthio)acetate (III).

Synthesis 4-((5-(alkylthio)-4-phenyl-4*H*-1,2,4-triazole-3-yl)methyl)-morpholine (IV) and (V). To a stirred solution of triazole (II) (0.01 mole) and triethylamine (0.01 mole) in absolute ethanol 20 ml, alkyl halide (0.01 mole) namely methyl iodide and / or butyl bromide was added. The reaction mixture was heated under reflux for 6 hrs. the solvent was removed under vacuum and the solid product was filtered off and crystallized from proper solvent to give white crystals of (IV) and (V) respectively.

Synthesis (3-(morpholinomethyl)-4-phenyl-5-thioxo-4,5-dihydro-1*H*-1,2,4-thiazol-1-yl)(phenyl)-methanone (VI). To a suspension of triazole (II) (0.01 mole) and triethylamine (0.01 mole) in dry benzene 20 ml, benzoyl chloride (0.01 mole) was added. The reaction mixture was heated under reflux for 6 hrs. The solid product obtained after cooling was filtered off and crystallized from chloroform to give white crystals of (3-(morpholinomethyl)-4-phenyl-5-thioxo-4,5-dihydro-1*H*-1,2,4-thiazol-1-yl)(phenyl)-methanone (VI).

REFERENCES

- [1] Kaldrikyan M.A., Grigoryan L.A., Melik-Ogandzhanyan R.G., Arsenyan F.G. *Synthesis and antitumor activity of some benzofuryl substituted 1,2,4-triazoles and triazolothiones-5*, CHC, **2011**, 4, 15-17 (in Russ.).
- [2] Joule J.A., Mills K. *Heterocyclic Chemistry*, 4th ed., Blackwell Science, Oxford, **2002**. 589 pp. (in Eng).
- [3] Sainova G.A., Tolisbaev E.B., Zharmenova M.B., *The search of new pharmacologically active substances in a series of piperidine derivatives*, International Research Journal: A Collection of the results of XVIII International Research Journal Conference, **2013**, Vol. 4, 8, 43-44 (in Russ.).
- [4] Mashkovskiy M.D. *Drugs*. 16th ed. M.: Novaya volna, **2012**. 1432 p. (in Russ.).
- [5] Vartanyan R.S. *Synthesis of Essential Medicines*. M.: MIA, **2005**. 845 p. (in Russ.).
- [6] Soldatenkov A.T., Koljadina N.M., Shendrik I.V. *Fundamentals of organic chemistry of medicinal substances*. M.: Himija, **2001**. 192 p. (in Russ.).
- [7] Bordyushkov Yu.N., Golotina L.Yu. *Rihlokain as an immunocorrector and activator of antitumor drugs*, Kuban Research Medical Bulletin, **2000**, 4, 107-109 (in Russ.).
- [8] Semina I.I., Shilovskaya E.V., Tihonova N.A., Baychurina A.Z., Tarasova R.I., Garaev R.S. *Anticonvulsant activity of unsubstituted 1,2,4-triazoles*, Him.-farm. zh., **2010**, Vol. 36, 2, 3-5 (in Russ.).
- [9] Dyusebaeva M.A., Kalugin S.N., Ahmedova Sh.S. *Synthesis and intramolecular cyclization of thiosemicarbazide morpholilacetic acid*, Chemical Bulletin of KazNU, **2015**, 2 (in Russ.).

ЛИТЕРАТУРА

- [1] Калдрикийн М.А., Григорян Л.А., Мелик-Оганджян Р.Г., Арсенян Ф.Г. Синтез и противоопухолевая активность некоторых бензофурилзамещенных 1,2,4-триазолов и триазолинтхионов-5 // ХГС. – 2011. – № 4. – Т. 45. – С. 15-17.
- [2] Joule J.A., Mills K. *Heterocyclic Chemistry*, 4th ed., Blackwell Science, Oxford, 2002. 589 pp.
- [3] Саинова Г.А., Толисбаев Е.Б., Жарменова М.Б. Поиск новых фармакологически активных веществ в ряду производных пиперидина // Международный научно-исследовательский журнал: Сборник по результатам International Research Journal Conference XVIII. Екатеринбург. : МНИЖ – 2013. – № 8. – Ч. 4. – С. 43-44.
- [4] Машковский М.Д. Лекарственные средства. Издание шестнадцатое. – М.: Новая волна, 2012. – 1432 с.
- [5] Вартанян Р.С. Синтез основных лекарственных средств. – М. : МИА, 2005. – 845 с.
- [6] Солдатенков А.Т., Колядина Н.М., Шендрик И.В. Основы органической химии лекарственных веществ. – М.: Химия, 2001. – 192 с.
- [7] Бордюшков Ю.Н., Голотина Л.Ю. Рихлокаин как иммунокорректор и активатор противоопухолевых препаратов // Кубанский научный медицинский вестник. – 2000. – № 4. – С. 107-109.
- [8] Семина И.И., Шиловская Е.В., Тихонова Н.А., Байчурина А.З., Тарасова Р.И., Гараев Р.С. Противосудорожная активность незамещенных производных 1,2,4-триазолов // Хим.-фарм. ж. - 2010. - Т. 36. - № 2. - С. 3-5.
- [9] Дюсебаева М.А., Калугин С.Н., Ахмедова Ш.С. Синтез и внутримолекулярная циклизация тиосемикарбазида морфолилуксусной кислоты // Вестник КазНУ, Серия Химическая. – 2015. – №2. – С.

3-(МОРФОЛИНОМЕТИЛ)-4-ФЕНИЛ-1H-1,2,4-ТРИАЗОЛ-5(4H)-ТИОННЫҢ АЛКИЛДЕНУ РЕАКЦИЯЛАРЫ**М. А. Дюсебаева, Ж. Женис, Л. Б. Жаймухамбетова, Ш. С. Ахметодова**

Әл-Фараби атындағы Қазақ ұлттық университеті, Алматы, Қазақстан

Тірек сөздер: морфолин, тиосемикарбазид, триазол, циклдену.

Аннотация. 1,2,4-Триазол-5-тиондардың туындылары тион-тиолды топтасуы бар гетероциклді қосылыстарға жатады. Бұл 3-(морфолинометил)-4-фенил-1H-1,2,4-триазол-5(4H)-тионның (II) екі таутомерлі формада (IIa и IIb) кездесетіндігін білдіреді және осы қасиеті реакцияларды екі реакциялық орталықтар арқылы жүргізуге мүмкіндік береді (NH немесе SH топтар арқылы). 1,2,4-Триазол сақинасында орынбасарлардың болуы 1,2,4-триазол ядросында орынбасу реакцияларының жүруіне алып келеді. Орынбасқан 1,2,4-триазол-5(4H)-тионды (II) алкилдеген немесе ацилдеген кезде енуші топ азот атомына немесе күкірт атомына қосыла алады. Осылайша бромсірке қышқылының этил эфирімен, метилиодид және бромды бутилмен сілтілік ортада алкилдегенде, сәйкесінше S-алкилденген туындылар (III), (IV) және (V) синтезделді. Ал 1,2,4-триазол-5(4H)-тионды (II) хлорлы бензоилмен триэтиламин қатысында ацилдеу, нәтижесінде N-ацилденген туындыны (VI) берді. Алынған қосылыстардың құрылыстары элементтік талдау, ИК және Н-ЯМР спектрлермен дәлелденді.

**РЕАКЦИИ АЛКИЛИРОВАНИЯ
3-(МОРФОЛИНОМЕТИЛ)-4-ФЕНИЛ-1H-1,2,4-ТРИАЗОЛ-5(4H)-ТИОНА****М. А. Дюсебаева, Ж. Женис, Л. Б. Жаймухамбетова, Ш. С. Ахметодова**

Казахский национальный университет им. аль-Фараби, Алматы, Казахстан

Ключевые слова: морфолин, тиосемикарбазид, триазол, циклизация, S-алкилирование, N-ацилирование.

Аннотация. Производные 1,2,4-триазол-5-тионов относятся к гетероциклическим соединениям с тион-тиольной группировкой. Это предполагает таутомерные превращения 1,2,4-триазолин-5-тиона в 1,2,4-триазол-5-тиол. 3-(Морфолинометил)-4-фенил-1H-1,2,4-триазол-5(4H)-тион (II) существует в двух таутомерных формах (IIa и IIb). И это позволяет проводить реакции по двум реакционным центрам (по NH или SH группам). Наличие заместителей в кольце 1,2,4-триазола способствует неоднозначному протеканию реакции замещения в ядре 1,2,4-триазола. При алкилировании или ацилировании замещенного 1,2,4-триазол-5(4H)-тиона (II) вступающая группа может присоединиться либо к атому азота, либо к атому серы. Таким образом, при алкилировании этиловым эфиром бромуксусной кислоты, метилиодидом и бромистым бутилом в щелочной среде синтезированы S-алкилированные производные (III), (IV) и (V) соответственно. В то время как ацилирование 1,2,4-триазол-5(4H)-тиона (II) хлористым бензоилом в присутствии триэтиламина привело к N-ацилированному производному (VI). Структуры полученных соединений были подтверждены данными элементного анализа, ИК и спектров Н-ЯМР.

Поступила 03.12.2015г.

Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

www.nauka-nanrk.kz

<http://www.chemistry-technology.kz/index.php/ru/>

Редакторы: *М. С. Ахметова, Т. А. Апендиев*
Верстка на компьютере *Д. Н. Калкабековой*

Подписано в печать 18.12.2015.

Формат 60x881/8. Бумага офсетная. Печать – ризограф.

8,5 п.л. Тираж 300. Заказ 6.