

ISSN 2518-1491 (Online),  
ISSN 2224-5286 (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ  
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ

# Х А Б А Р Л А Р Ы

**ИЗВЕСТИЯ**

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК  
РЕСПУБЛИКИ КАЗАХСТАН

**N E W S**

OF THE ACADEMY OF SCIENCES  
OF THE REPUBLIC OF KAZAKHSTAN

**ХИМИЯ ЖӘНЕ ТЕХНОЛОГИЯ  
СЕРИЯСЫ**

◆  
**СЕРИЯ**  
**ХИМИИ И ТЕХНОЛОГИИ**  
◆  
**SERIES**  
**CHEMISTRY AND TECHNOLOGY**

**1 (427)**

**ҚАҢТАР – АҚПАН 2018 ж.  
ЯНВАРЬ – ФЕВРАЛЬ 2018 г.  
JANUARY – FEBRUARY 2018**

1947 ЖЫЛДЫҢ ҚАҢТАР АЙЫНАН ШЫҒА БАСТАҒАН  
ИЗДАЕТСЯ С ЯНВАРЯ 1947 ГОДА  
PUBLISHED SINCE JANUARY 1947

**ЖЫЛЫНА 6 РЕТ ШЫҒАДЫ  
ВЫХОДИТ 6 РАЗ В ГОД  
PUBLISHED 6 TIMES A YEAR**

АЛМАТЫ, ҚР ҰФА  
АЛМАТЫ, НАН РК  
ALMATY, NAS RK

*NAS RK is pleased to announce that News of NAS RK. Series of chemistry and technologies scientific journal has been accepted for indexing in the Emerging Sources Citation Index, a new edition of Web of Science. Content in this index is under consideration by Clarivate Analytics to be accepted in the Science Citation Index Expanded, the Social Sciences Citation Index, and the Arts & Humanities Citation Index. The quality and depth of content Web of Science offers to researchers, authors, publishers, and institutions sets it apart from other research databases. The inclusion of News of NAS RK. Series of chemistry and technologies in the Emerging Sources Citation Index demonstrates our dedication to providing the most relevant and influential content of chemical sciences to our community.*

Қазақстан Республикасы Ұлттық ғылым академиясы "ҚР ҰҒА Хабарлары. Химия және технология сериясы" ғылыми журналының Web of Science-тің жаңаланған нұсқасы Emerging Sources Citation Index-те индекстелуге қабылданғанын хабарлайды. Бұл индекстелу барысында Clarivate Analytics компаниясы журналды одан әрі the Science Citation Index Expanded, the Social Sciences Citation Index және the Arts & Humanities Citation Index-ке қабылдау мәселесін қарастыруды. Web of Science зерттеушілер, авторлар, баспашилар мен мекемелерге контент тереңдігі мен сапасын ұсынады. ҚР ҰҒА Хабарлары. Химия және технология сериясы Emerging Sources Citation Index-ке енүі біздің қоғамдастық үшін ең өзекті және беделді химиялық ғылымдар бойынша контентке адалдығымызды білдіреді.

НАН РК сообщает, что научный журнал «Известия НАН РК. Серия химии и технологий» был принят для индексирования в Emerging Sources Citation Index, обновленной версии Web of Science. Содержание в этом индексировании находится в стадии рассмотрения компанией Clarivate Analytics для дальнейшего принятия журнала в the Science Citation Index Expanded, the Social Sciences Citation Index и the Arts & Humanities Citation Index. Web of Science предлагает качество и глубину контента для исследователей, авторов, издателей и учреждений. Включение Известия НАН РК в Emerging Sources Citation Index демонстрирует нашу приверженность к наиболее актуальному и влиятельному контенту по химическим наукам для нашего сообщества.

**Бас редакторы**  
х.ғ.д., проф., ҚР ҮҒА академигі **М.Ж. Жұрынов**

Редакция алқасы:

**Ағабеков В.Е.** проф., академик (Белорус)  
**Волков С.В.** проф., академик (Украина)  
**Воротынцев М.А.** проф., академик (Ресей)  
**Газалиев А.М.** проф., академик (Қазақстан)  
**Ергожин Е.Е.** проф., академик (Қазақстан)  
**Жармағамбетова А.К.** проф. (Қазақстан), бас ред. орынбасары  
**Жоробекова Ш.Ж.** проф., академик (Қырғыстан)  
**Итқұлова Ш.С.** проф. (Қазақстан)  
**Манташян А.А.** проф., академик (Армения)  
**Пралиев К.Д.** проф., академик (Қазақстан)  
**Баешов А.Б.** проф., академик (Қазақстан)  
**Бұркітбаев М.М.** проф., академик (Қазақстан)  
**Джусипбеков У.Ж.** проф. корр.-мүшесі (Қазақстан)  
**Молдахметов М.З.** проф., академик (Қазақстан)  
**Мансуров З.А.** проф. (Қазақстан)  
**Наурызбаев М.К.** проф. (Қазақстан)  
**Рудик В.** проф., академик (Молдова)  
**Рахимов К.Д.** проф. академик (Қазақстан)  
**Стрельцов Е.** проф. (Белорус)  
**Тәшімов Л.Т.** проф., академик (Қазақстан)  
**Тодераш И.** проф., академик (Молдова)  
**Халиков Д.Х.** проф., академик (Тәжікстан)  
**Фарзалиев В.** проф., академик (Әзірбайжан)

**«ҚР ҮҒА Хабарлары. Химия және технология сериясы».**

**ISSN 2518-1491 (Online),**

**ISSN 2224-5286 (Print)**

Меншіктенуші: «Қазақстан Республикасының Үлттық ғылым академиясы» Республикалық қоғамдық бірлестігі (Алматы қ.)

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде 30.04.2010 ж. берілген №1089-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік

Мерзімділігі: жылдан 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,  
[www.nauka-nanrk.kz](http://www.nauka-nanrk.kz) / [chemistry-technology.kz](http://chemistry-technology.kz)

---

© Қазақстан Республикасының Үлттық ғылым академиясы, 2018

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Г л а в н ы й р е д а к т о р  
д.х.н., проф., академик НАН РК **М. Ж. Журинов**

Р е д а к ц и о н на я к о л л е г и я:

**Агабеков В.Е.** проф., академик (Беларусь)  
**Волков С.В.** проф., академик (Украина)  
**Воротынцев М.А.** проф., академик (Россия)  
**Газалиев А.М.** проф., академик (Казахстан)  
**Ергожин Е.Е.** проф., академик (Казахстан)  
**Жармагамбетова А.К.** проф. (Казахстан), зам. гл. ред.  
**Жоробекова Ш.Ж.** проф., академик (Кыргызстан)  
**Иткулова Ш.С.** проф. (Казахстан)  
**Манташян А.А.** проф., академик (Армения)  
**Пралиев К.Д.** проф., академик (Казахстан)  
**Баешов А.Б.** проф., академик (Казахстан)  
**Буркитбаев М.М.** проф., академик (Казахстан)  
**Джусипбеков У.Ж.** проф. чл.-корр. (Казахстан)  
**Мулдахметов М.З.** проф., академик (Казахстан)  
**Мансуров З.А.** проф. (Казахстан)  
**Наурызбаев М.К.** проф. (Казахстан)  
**Рудик В.** проф., академик (Молдова)  
**Рахимов К.Д.** проф. академик (Казахстан)  
**Стрельцов Е.** проф. (Беларусь)  
**Ташимов Л.Т.** проф., академик (Казахстан)  
**Тодераш И.** проф., академик (Молдова)  
**Халиков Д.Х.** проф., академик (Таджикистан)  
**Фарзалиев В.** проф., академик (Азербайджан)

**«Известия НАН РК. Серия химии и технологии».**

**ISSN 2518-1491 (Online),**

**ISSN 2224-5286 (Print)**

Собственник: Республиканское общественное объединение «Национальная академия наук Республики Казахстан» (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №10893-Ж, выданное 30.04.2010 г.

Периодичность: 6 раз в год

Тираж: 300 экземпляров

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел. 272-13-19, 272-13-18,  
<http://nauka-nanrk.kz> / [chemistry-technology.kz](http://chemistry-technology.kz)

---

© Национальная академия наук Республики Казахстан, 2018

Адрес редакции: 050100, г. Алматы, ул. Кунаева, 142,  
Институт органического катализа и электрохимии им. Д. В. Сокольского,  
каб. 310, тел. 291-62-80, факс 291-57-22, e-mail:[orgcat@nursat.kz](mailto:orgcat@nursat.kz)

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75

Editor in chief  
doctor of chemistry, professor, academician of NAS RK **M.Zh. Zhurinov**

Editorial board:

**Agabekov V.Ye.** prof., academician (Belarus)  
**Volkov S.V.** prof., academician (Ukraine)  
**Vorotyntsev M.A.** prof., academician (Russia)  
**Gazaliyev A.M.** prof., academician (Kazakhstan)  
**Yergozhin Ye.Ye.** prof., academician (Kazakhstan)  
**Zharmagambetova A.K.** prof. (Kazakhstan), deputy editor in chief  
**Zhorobekova Sh.Zh.** prof., academician (Kyrgyzstan)  
**Itkulova Sh.S.** prof. (Kazakhstan)  
**Mantashyan A.A.** prof., academician (Armenia)  
**Praliyev K.D.** prof., academician (Kazakhstan)  
**Bayeshov A.B.** prof., academician (Kazakhstan)  
**Burkitbayev M.M.** prof., academician (Kazakhstan)  
**Dzhusipbekov U.Zh.** prof., corr. member (Kazakhstan)  
**Muldakhmetov M.Z.** prof., academician (Kazakhstan)  
**Mansurov Z.A.** prof. (Kazakhstan)  
**Nauryzbayev M.K.** prof. (Kazakhstan)  
**Rudik V.** prof., academician (Moldova)  
**Rakhimov K.D.** prof., academician (Kazakhstan)  
**Streltsov Ye.** prof. (Belarus)  
**Tashimov L.T.** prof., academician (Kazakhstan)  
**Toderash I.** prof., academician (Moldova)  
**Khalikov D.Kh.** prof., academician (Tadzhikistan)  
**Farzaliyev V.** prof., academician (Azerbaijan)

**News of the National Academy of Sciences of the Republic of Kazakhstan. Series of chemistry and technology.**  
**ISSN 2518-1491 (Online),**  
**ISSN 2224-5286 (Print)**

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)  
The certificate of registration of a periodic printed publication in the Committee of Information and Archives of the Ministry of Culture and Information of the Republic of Kazakhstan N 10893-Ж, issued 30.04.2010

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,  
<http://nauka-nanrk.kz> / [chemistry-technology.kz](http://chemistry-technology.kz)

---

© National Academy of Sciences of the Republic of Kazakhstan, 2018

Editorial address: Institute of Organic Catalysis and Electrochemistry named after D. V. Sokolsky  
142, Kunayev str., of. 310, Almaty, 050100, tel. 291-62-80, fax 291-57-22,  
e-mail: [orgeat@nursat.kz](mailto:orgeat@nursat.kz)

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

**NEWS**

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

**SERIES CHEMISTRY AND TECHNOLOGY**

ISSN 2224-5286

Volume 1, Number 427 (2018), 53 – 58

**Zh. N. Komekbay, Z. B. Halmenova, A. K. Umbetova, A.G Bisenbay**

Kazakh national university named after al-Farabi, Almaty, Kazakhstan

E-mail: jaziko\_94\_21@mail.ru, zaure.halmenova@mail.ru, alma\_0875@mail.ru, aimanka.b-97@mail.ru

**PHYTOCHEMICAL ANALYSIS AND DEVELOPMENT  
OF PRODUCTION OF BIOLOGICALLY ACTIVE COMPLEX  
ON THE BASIS OF RAW MELISSA OFFICINALIS L**

**Abstract.** In this study, there was developed a comprehensive study of wild and cultured plants of the genus *Melissa* (*Melissa officinalis* L.). It was determined the purity of the raw materials: moisture, total ash, ash insoluble in 10% HCl, sulphate ash, extractives. Macro- and microelement composition of total ash by atomic absorption spectroscopy was analyzed. The analysis of the component composition of the main classes of natural substances was conducted. There were worked out the basic technological parameters of obtaining biologically active complex of the studied plant species by varying the nature of the extractant, its ratio of raw materials, time, and frequency extraction.

**Key words:** *Melissa officinalis* L., extractives, moisture content, total ash, ash insoluble in HCl, sulphate ash, macro - and microelement composition, atomic absorption spectroscopy, phytochemical analysis.

One of the priorities of the development of home science and practice of chemistry of natural compounds is the more complete use of own resources of wild and cultivated plant raw materials and the creation of effective drugs based on it, affordable and not inferior in quality to foreign analogues. Among natural biologically active compounds used for the treatment of upper respiratory tract diseases, special attention should be paid to plants of the family *Lamiaceae Lindl* (clear flow) [1].

The genus *Melissa* (*melissa*) includes from 2 to 10 species. It grows like a weed plant in orchards, roads, fields, occasionally wild in the lower belt of mountains, and this plant is also cultivated. The most valuable species is *Melissa officinalis* L (lemon balm), home to which is the eastern Mediterranean region. Cultivate melissa medicinal in many countries of the world, where it is registered as pharmacopoeial plants.

In the culture this plant is introduced throughout Europe and North America and in Kazakhstan the plant is introduced in the regions like South Kazakhstan (Shymkent), Zhambyl (Taraz), Kyzylorda, Almaty regions. Wild species of *Melissa officinalis* L are common in Central and Southern Europe, the Caucasus, Middle and Near Asia, North Africa and North America [2].

The biological value of the raw material of lemon balm is caused by a complex of biologically active substances, such as ether compounds, phenolic substances, vitamins.

Phenolic compounds of *Melissa officinalis* are represented by phenolcarboxylic acids and their derivatives, flavonoids and coumarins. Analysis of literature data shows that, in the lemon balm medicated grown in Europe, n-coumaric, ferulic, kaftaric and coffee acids have been identified. Other researchers have identified rosemary, coffee and protocatechuic acids.

In addition, flavonoids, the glycosides of luteolin and apigenin, are characteristic of this plant. The aqueous extract of lemon balm contains hydrolysable tannins in an amount of 4.32% and flavonoids in an amount of 2.06% [3].

Healing properties of the aerial part of the lemon balm are caused by the high content of essential oil. Its most characteristic components are monoterpenes - citral, geraniol, nerol, citronellol, citronellal. *Melissa* essential oil also contains linalool, geranilacetate, myrcene, n-cymene, beta-cariophyllene oxide, beta-cariophyllene and other terpenoids, and more than 200 compounds have been isolated and described in total.

The second group of components of essential oil are phenylpropanoids, among which the most characteristic is rosmarinic acid. Phenylpropanoids are a class of plant organic compounds of the aromatic series, which are synthesized in a shikimate way, mainly through the amino acid phenylalanine. A characteristic structural fragment is a benzene ring with a branched three-carbon chain attached to it. Phenylpropanoids have a wide range of functions - protection from herbivorous animals and microbial diseases, protection against ultraviolet light, serve as structural components of cellular stains, pigments, act as signal molecules. Phenylpropanoids are also represented by ethyl ester of rosmarinic acid, caffeic acid, chlorogenic acid, n-coumaric acid, ferulic acid and synapic acids. The content of rosmarinic acid in melissa leaves is from 0.54 to 1.79% [4].

Melissa leaves also contain triterpenes - ursolic and oleanolic acids (0.50% and 0.17%, respectively) and their derivatives, terpenoids - nerol, geraniol, nerolic acid glucosides. They are found bitterness, coumarins (esketuin), up to 5% tannins, succinic acid, mucus, tetrasaccharide stachyose (a combination of two galactose residues with glucose and fructose), carotene (0.007-0.01%), vitamins C (0.15 %), B1, B2, E [1, 2].

Melissa is widely used in medicine, in the perfumery, cosmetics and food industries in many countries. Raw melissa has sedative, spasmolytic, immunomodulating, antidepressant, antihistamine, antioxidant, anti-inflammatory and antimicrobial effects. In addition, it was found that this plant has antiviral activity against viral infections, such as smallpox, influenza, herpes [5].

Medicines, which include lemon balm, have pronounced soothing, antispasmodic and carminative properties. It has been established that melissa shows an easy hypnotic effect. Such pharmacological activity is mainly due to the components of the essential oil. Sedative and spasmolytic effects are manifested with the application of small doses of lemon balm, and the subsequent increase does not enhance these effects [6].

In the seeds of lemon balm it contains up to 20% of fatty oil as it contains up to 20% of fatty oil.

Melissa tincture shows protective effect with experimental stomach ulcer. At the same time, it is established that it enhances the motility of the stomach, has choleric and haemostatic properties. In the experimental animals, the antispasmodic effect of melissa has been established. Its tincture reduces the tension of the smooth muscles of the intestine, shows bronchodilator properties.

Melissa essential oils show anti-inflammatory, bacteriostatic and antiviral properties. Japanese scientists conducted a study of the antimicrobial activity of essential oil components of the plant *Melissa officinalis L* against a number of pathogenic fungi and microbacteria of tuberculosis. The most active were aldehydes (citral, citronellal), and less active alcohols (geraniol) essential oil of the plant [7].

**The purpose** of the research work is to substantiate the possibility of using the cultivated and wild-growing species *Melissa officinalis L*, introduced under the conditions of the Almaty region to obtain an extract with the subsequent study of the chemical composition.

**The objects** of the study are samples of a plant of the family *Lamiaceae* genus *Melissa* (*Melissa*) and its appearance *Melissa officinalis L* (*Melissa officinalis*). Raw materials were individually cultivated at the experimental site of the laboratory of medicinal plants of the institute of phytointroduction and botany under the Ministry of Science and Education of the Republic of Kazakhstan, the city of Almaty and the wild type of lemon balm prepared in Almaty region.

### Experimental part and discussion

The selection of raw material is determined by its former prisms and personal indications (to exert, to keep, to contain explosive substances).

All the indications of the distribution were determined by the methods of the SP RK, the Euphemistic Map and other literature sources [8, 9]. The data are presented in Table 1.

Table 1 – Numerical requirement for publication of the type of *Melissa officinalis L*

References are requested	Plant <i>Melissa officinalis L</i>	
	Cultured	Wild-growing
Implication, %	5,36	4,42
Total wt%	7,91	10,60
Sulphate Salt, %	15,36	15,74
HCl of ash, %	13,80	14,68
Containment of detective substances (70% alcohol)	41,40	38,03

As can be seen from the data presented in Table 1, the resistance of the plant in a plant of the cultivated species (5.36%) is greater than that of the plant in the sample of the wild-growing species (4.42%) less.

The content of extractives in medicinal plant raw materials is an important numerical indicator that determines its good quality, especially for those types of raw materials in which the quantitative determination of active substances is not carried out.

Depending on the chemical composition of the medicinal plant material and the solvent used, certain active and concomitant substances pass into the extraction.

The solvent, which should be taken when determining extractive substances, is indicated in the relevant specification for this type of raw material. Usually it is the same solvent that is used when preparing a tincture or extract from this raw material. Most often it is ethyl alcohol (50 or 70%) or water.

From the data of Table 1 it follows that the greatest amount of extractive substances is extracted by the cultivated plant species.

In the plant raw material, the ash of total, sulfate ash, ashes are insoluble in 10% HCl, which is the residue after treatment with the total HCl ash and consists mainly of silicates, which are a natural component for some objects, but more often the result of contamination of the raw materials with sand, earth and stones. Thus, the increased content of insoluble in hydrochloric acid part of the ash indicates a significant content in the vegetable raw material of the mineral impurity. The amount of sulfate ash is commensurate with the content of metals in plants that form sulfate-insoluble in water. The content of all ash species in the aerial part of the plant does not exceed the maximum acceptable value for pharmacopoeial samples.

The next parameter of the definition is the mineral composition. The investigated plant species have a high ash content.

Mineral elements by their content in the plant are divided into macroelements, microelements and ultramicroelements. The macroelements include Na, K, Ca, Mg, their content in the ash is measured in hundredths of a percent. Microelements: Zn, Cu, Ni, Mn, Fe.

The aboveground part contains macro elements: K, Ca, Na, Mg; microelements: Mn, Fe, Cu, Zn, Ni. *Potassium and sodium* play a leading role in regulating the water-salt balance and acid-base balance of the body. *Calcium* plays a huge role in the life of the human body. The human body contains 1000-1200 g of calcium, 99% - is included in bone tissue, dentin, enamel of the teeth, and 1% plays an extremely important role as intracellular calcium, blood calcium and tissue fluid, that is, it plays an important role in the formation of bones. *Magnesium* participates in many processes occurring in the body - in energy production, glucose assimilation, neurotransmission, protein synthesis, bone tissue building, regulation of relaxation and tension of blood vessels and muscles. *Manganese* affects the development of the skeleton, participating in the process of osteogenesis, and therefore is necessary for normal growth. Manganese participates in the reactions of immunity, in blood and tissue respiration, supports reproductive functions, participates in the regulation of carbohydrate and lipid metabolism. *Zinc* is part of the active center of several hundred metal-enzymes. It is necessary for the functioning of DNA and RNA polymerases, which control the processes of transmission of hereditary information and the biosynthesis of proteins, and thereby the reparative processes in the body. *Nickel* is involved in stimulating the processes of hematopoiesis, the activation of certain enzymes. It has a high ability to enhance redox processes in tissues. Nickel in combination with cobalt, iron, copper participates in the processes of hematopoiesis, and independently - in the exchange of fats, providing cells with oxygen. In certain doses, it activates the action of insulin. *Iron* is the most important trace element, takes part in respiration, hematopoiesis, immunobiological and oxidation-reduction reactions, is part of more than 100 enzymes [10, 11].

In the total ash, the content of macro- and microelements was determined by atomic absorption spectroscopy. The data are presented in Table 2-3.

From the data of Tables 2 and 3, it should be noted that the greatest amount of macro- and microelements is in the plant *Melissa officinalis L*, a wild-growing species. In the above-mentioned mass of the cultivated and wild-growing species, the predominant micelles are Fe. In the cultivated form of the plant, an increased content of macroelements as K, Na, Ca and in the wild form Mg and K is noted. The content of heavy metals does not exceed the maximum permissible norms [11].

Table 2 - Macroelements content - K, Ca, Na, Mg

Macroelements	Plant <i>Melissa officinalis L</i>	
	Cultured, %	Wild-growing, %
K	$1.173 \cdot 10^{-3}$	$0.737 \cdot 10^{-3}$
Na	$0.802 \cdot 10^{-3}$	$0.221 \cdot 10^{-3}$
Ca	$0.639 \cdot 10^{-3}$	$0.178 \cdot 10^{-3}$
Mg	$0.313 \cdot 10^{-3}$	$1.401 \cdot 10^{-3}$

Table 3 - Micronutrients content of Fe, Zn, Mn, Cu, Ni

Microelements	Plant <i>Melissa officinalis L</i>	
	Cultured, %	Wild-growing, %
Cu	$0.716 \cdot 10^{-3}$	$0.394 \cdot 10^{-3}$
Fe	$4.387 \cdot 10^{-3}$	$1.266 \cdot 10^{-3}$
Mn	$0.361 \cdot 10^{-3}$	$0.677 \cdot 10^{-3}$
Ni	0	$0.502 \cdot 10^{-3}$
Zn	$0.486 \cdot 10^{-3}$	$0.335 \cdot 10^{-3}$

A comparative phytochemical analysis of the above-ground mass of the plant on the main classes of biologically active substances was carried out. The data are presented in Table 4.

Table 4 - Phytochemical analysis of the distribution of *Melissa officinalis L* cultivated and wild-growing species

BAS	Developers	<i>Melissa officinalis L</i>	
		Cultured	Wild-growing
Carbohydrates	o-toloidin	green	green
Tannins	ZHAK	blue	blue
	FeCl <sub>3</sub>	blue	blue
Flavonoids	NH <sub>3</sub>	yellow	yellow
	AlCl <sub>3</sub>	bright yellow	bright yellow
	SiHNO	orange-red	orange-red
Cataracts	KMnO <sub>4</sub>	bleaching	bleaching
Alkaloids	Phosphoric acid	bleaching	bleaching
Amino acids	ninhydrin	purple	purple
Carboxylic acids	urea	brown	brown
	MgAc <sub>2</sub>	-	-

Through phytochemical analysis using diagnostic gums in the aerial part of a plant of the type *Melissa officinalis L*, there were discovered the main groups of biological active substance such as tannins, amino acids, alkaloids, phenolic compounds, organic acids, flavonoids, carotenoids [12].

Method of paper chromatography in the use of reliable samples in the following types of pockets identified carbohydrates and amino acids.

The optimal technology for isolating the extract from the melissa plant was developed taking into account the requirements of the SP RK for the processing of plant raw materials [13, 14].

An important parameter in the technology of obtaining a plant extract is the ratio of raw materials and solvent from 1: 4 to 1: 8. Over 5 g of the aboveground part was extracted with different volumes of 50%, 70%, 90% ethyl alcohol. The constant factors of the extraction process were: extraction time (24 hours) and temperature (23-25 °C). These parameters for obtaining the plant extract are shown in Table 5.

From these tables it follows that the optimal extractant was 70% ethanol. The greatest yield of the percentage of the extract shows extraction with 70% ethanol at a feed: extractant ratio of 1: 8, the amount of dry extract in cultivated form was 7.94%, and in the wild 10.264%.

Another important parameter in the technology of extracting extracts is the ratio of the selected extract with the custome. To determine the optimum volume, the selected extractor changes the content of

the raw material and the filter from 1: 5 to 1: 8. To 5 g of the nasis part of the cultivated and wild-growing species *Melissa officinalis L* 70% ethyl alcohol. With this constant factor, the process of extraction was: time of extraction (24 hours) and temperature (23-25 ° C). The data are presented in Table 6.

Table 5 - Determination of the optimum extractant for the extraction of the raw materials

m,r const	Solvents	t,hour const	T,°C const	m(r):v (ml)	The amount of dry extract, % <i>Melissa officinalis L</i>	
					Cultured	Wild-growing
5	50% ethanol	24	23-25°C	1:4	0,528	0,783
				1:6	2,501	2,670
				1:8	5,124	8,574
5	70% ethanol	24	23-25°C	1:4	2,604	3,211
				1:6	3,25	4,454
				1:8	7,94	10,264
5	90% ethanol	24	23-25°C	1:4	0,447	0,695
				1:6	1,121	1,880
				1:8	3,344	3,212

Table 6 - Determination of the optimum ratio of raw material and extractant

Mass of raw materials, g const		5	5	5
Extraction time, hour, const		24	24	24
Extraction temperature, °C const		23-25°C	23-25°C	23-25°C
Ratio of raw materials (g) and extractant (ml)		1:4	1:6	1:8
V2 volume of the filtered extract, ml	Cultured	3,5	5,0	9,5
	Wild-growing	4,0	6,7	11,3
Extraction amount, %	Cultured	1,76	3,25	5,24
	Wild-growing	2,24	4,62	6,89

With the selected extract (70% ethanol), the optimum value of the extract 1:8 is shown in cultivated form, the amount of the extract is 5.24%, and in the dicorbic form 6.89%; at a temperature of 24-25 ° C and a time of 24 hours.

The accuracy of the parameters of the parameters "raw-extractor" is determined primarily by economic measures, as for a business, but the amount of the used extractor is significant.

Determination of the extraction time is an important parameter, so it is necessary to determine when the entire complex of biological active substances is recovered. The influence of the extraction time of the raw material on the yield of the extract was studied. Extraction was carried out with 70% ethanol at a raw material: extractant ratio of 1: 8. The data are presented in Table 7.

Table 7 - Determination of extraction time

Extractant (ml), const		70% ethanol	70 % ethanol	70 %ethanol
Ratio of raw materials (g) and extractant (ml), const		1:8	1:8	1:8
Time, (hour)		24	48	72
V2 volume of the filtered extract, ml	Cultured	17	2	7
	Wild-growing	13	8	11
Extraction amount, %	Cultured	0.3939	0.0605	0.1365
	Wild-growing	0.2902	0.2785	0.2336

On the basis of these arguments and the data obtained, the following regime was proved to be optimal: extraction of 70% by ethanol in 24 hours at a temperature of no more than 23-25 ° C with raw materials: extractant ratio 1: 8. Under this regime, the amount of extract was 0.3939% cultivated, 0.2902% in the wild.

**Acknowledgement.** According to the results of the research work, a comparative analysis of the chemical composition of the above-ground part of the cultivated and wild-growing species of the family

*Melissa (Melissa officinalis L)* of the family *Lamiaceae* was performed; Technological parameters were worked out: different concentrations of extractants; dependence of the ratio of raw materials - solvent; the process dependence on time and the number of extractions. The optimum condition for obtaining a plant substance is an extractant of 70% ethyl alcohol, the ratio of extractant and raw material is 1: 8, extraction time is 24 hours, temperature is 23-25 ° C.

#### REFERENCES

- [1] Flopa Kazaxctana. Cemejctva *Melissa Lamiaceae Lindl.* T.6. Alma-Ata, **1963** с. 437-438. (in Russ.)
- [2] Zuzuk B.M. *Melissa lekarstvennaja (Melissa officinalis L.)* B.M. Zuzuk, R.V. Kucik. Provizor. **2002**. № 1. S. 36-39. (in Russ.)
- [3] Patora J. *Flavonoids from lemon balm (Melissa officinalis L., Lamiaceae)* J. Patora, B. Klimek: Acta Polonica Pharmaceutica. **2002**. Vol. 59, № 2. P. 139-143. (in Eng.)
- [4] *Evaluation of phenolic acid derivatives and essential oil content in some Melissa officinalis L. varieties* Oniga, L. Vlase, A. Toiu [et al.] Farmacia. **2010**, Vol. 58, № 6.P. 764-769. (in Eng.)
- [5] *Antiherpes effect of Melissa officinalis L. extracts* Z. Dimitrova, N. Manolova, S. Pancheva [et al.] Acta Microbiol Bulg. **1993**. Vol. 29. R. 65-72
- [6] *Antioxidant activity of Melissa officinalis leaves*. E. Koksal, E. Bursal, E. Dikici [et al.] J. Med. Plant.Res. **2011**. Vol. 5, № 2. P. 217- 222. (in Eng.)
- [7] Mupav'eva D.A, Cal'mina L.A, Jakovlev G.P. *Fapmakognozija*. M.: Med., **2002**. 656 c. (in Russ.)
- [8] *Gocudapctvennaja fapmakopeja Pecpubliki Kazaxstan*. T. 1. Almaty: Izd. dom «Zhibek Zholy», **2008**. 592 c. (in Russ.)
- [9] European Pharmacopoeia. Strasburg, **2001**. P. 1705. (in Eng.)
- [10] Dobrynina N.A. *Biologicheskaja rol' nekotoryh himicheskikh jelementov*. Himija v shkole. **1991**. №2. S. 6-14.(in Russ.)
- [11] Miftahova A.F. *Fitohimicheskoe izuchenie rastenij nekotoryh vidov semejstva jasnotkovyh*. Avtoref. dis. kan-ta. him. nauk. Almaty, **2002**. 23-25 S. (in Russ.)
- [12] *Xromatografija na bumage pod red.* Xajsa I.M., Maceka M. M.: «Nacional'naja literatura», **1962**. 852 s. (in Russ.)
- [13] P.A.Muzychkina, D.Ju.Kopul'kin, Zh.A.Abilov *Kachectvennyj i kolichectvennyj analiz osonovnyx grupp BAV v lekapctvennom pachtel'nom cyp'e i fitopreparatax*. Almaty: Qazaq univepciteti, **2004**. 288c(in Russ.)
- [14] S.A. Minina, I.E. Kauhova. *Himija i tehnologija fitopreparatov*. M.: **2004** g.(in Russ.)

**Ж.Н. Қемекбай, З.Б. Халменова, А.К. Үмбетова, А.Ғ Бисенбай**

әл-Фараби атындағы Қазақ ұлттық университеті, Алматы, Қазақстан

#### ***MELISSA OFFICINALIS L ӨСІМДІГІ НЕГІЗІНДЕ ФИТОХИМИЯЛЫҚ АНАЛИЗ ЖАСАУ ЖӘНЕ БИОЛОГИЯЛЫҚ БЕЛСЕНДІ КЕШЕНДІ АЛУ***

**Аннотация.** Бұл зерттеуде *Melissa (Melissa officinalis L)* тұқымдасына жататын есімдіктің мәдени және жабайы түрлерінің кешенді негізі жетілдірілді. Шикізат сапалылығы: ылғалдылық, жалпы күлділік, HCl-да ерімейтін күлділік, сульфатты күлділік, экстрактивті заттар анықталды. Атомдық-абсорбциялық спектроскопиялық әдіспен жалпы күлділіліктің макро- және микроэлементтік құрамы талданды. Талдау компоненттік құрамның негізгі кластары табиги заттар. Зерттелініп отырған есімдік түрлерінен экстрагенттің табигатын, оның шикізатпен қатынасын, экстрактилеу уақыты мен жиілігін өзгерте отырып, биологиялық белсенді кешенді алу технологиясы өндедлі.

**Түйін сөздер:** *Melissa officinalis L*, экстрактивті заттар, ылғалдылық, жалпы күлділік, HCl да ерімейтін күлділік, сульфатты күлділік, макро- және микроэлементтік құрамы, атомдық-абсорбциялық спектроскопия, фитохимиялық анализ.

**МАЗМУНЫ**

<i>Ергөжин Е.Е., Бектенов Н.Ә., СенГүпта Арут К., Байдулаева А.Қ., Садыков Қ.А., Әбдіралиева Г.Е., Қалмуратова К.М., Рыспаева С.Б.</i> Эпоксиакрилат пен комплексондар негізіндегі жаңа комплекстүзгіш ион алмастырыштар арқылы стронций иондарын сорбциялау (ағылшын тілінде).....	6
<i>Ауелханкызы М., Славинская Н.А., Шабанова Т.А., Мансуров З.А.</i> Алленнің тотығуын және пиролизін модельдік зерттеу (ағылшын тілінде).....	12
<i>Ешіжанов А.А., Волненко А.А., Левданский А.Э., Корганбаев Б.Н.</i> Құрамалы туракты - қалықтамалы саптаманың эквивалентті диаметрін есептеуге (ағылшын тілінде).....	20
<i>Жұмадуллаев Д.К., Ешіжанов А.А., Волненко А.А., Левданский А.Э.</i> Беттік және араластырыштық жылу алмастырыштардың құбырлы шоғырындағы гидравликалық кедергіні есептеудің бірінғай тәсілдемесі (ағылшын тілінде) .....	25
<i>Савденбекова Б.Е., Оспанова А.К., Уваров Н.Ф.</i> Инженерлік технологияда белгілі бір мақсатты қасиетке ие перспективалық композитті материалдарды алу үшін мультижинақтау әдісін (LBL) қолдану (ағылшын тілінде).....	31
<i>Альчинбаева О., Сарбаева Қ.</i> Адам ағзасына химиялық ағартқыш заттарының әсері (ағылшын тілінде).....	38
<i>Үмбетова А.К., Слан Г.О., Омарова А.Т., Бурашева Г.Ш., Абидкулова К.Т.</i> Алматы өніріндегі <i>Atrapaxis virgata</i> өсімдігінің химиялық құрамын зерттеу (ағылшын тілінде).....	42
<i>Баешов Ә.Б., Нұрділаева Р.Н., Ташикенбаева Н.Ж., Өзлер М.Ә.</i> Айнымалы токпен поляризацияланған tot баспайтын болаттың еруі (ағылшын тілінде).....	46
<i>Көмекбай Ж.Н., Халменова З.Б., Үмбетова А.К., Бисенбай А.Ф.</i> <i>Melissa officinalis L</i> өсімдігі негізінде фитохимиялық анализ жасау және биологиялық белсенді кешен алу (ағылшын тілінде).....	53
<i>Мамырбекова А., Баешов А.Б., Касымова М.К., Мамырбекова А.</i> Микроорганизмдердің өсуіне газ тасымалдау қызметі бар перфтордекалиннің әсері (ағылшын тілінде).....	59
<i>Еспанова И.Д., Жусупова Л.А., Тапалова А.С., Анпазов Н.О.</i> Гексен-1 мен бутан қышқылының қосылу реакциясын микротолқындық белсендіріу (ағылшын тілінде).....	63
<i>Надиров Н.К., Некрасов В.Г., Солодова Е.В., Срымов Т., Суханбердиева Д.Т., Култоре М.А.</i> Жаңа буын жылышайы (ағылшын тілінде).....	70

\* \* \*

<i>Ергөжин Е.Е., Бектенов Н.Ә., СенГүпта Арут К., Байдулаева А.Қ., Садыков Қ.А., Әбдіралиева Г.Е., Қалмуратова К.М., Рыспаева С.Б.</i> Эпоксиакрилат пен комплексондар негізіндегі жаңа комплекстүзгіш ион алмастырыштар арқылы стронций иондарын сорбциялау (орыс тілінде).....	81
<i>Ешіжанов А.А., Волненко А.А., Левданский А.Э., Корганбаев Б.Н.</i> Құрамалы туракты - қалықтамалы саптаманың эквивалентті диаметрін есептеуге (орыс тілінде).....	87
<i>Жұмадуллаев Д.К., Ешіжанов А.А., Волненко А.А., Левданский А.Э.</i> Беттік және араластырыштық жылу алмастырыштардың құбырлы шоғырындағы гидравликалық кедергіні есептеудің бірінғай тәсілдемесі (орыс тілінде).....	93
<i>Савденбекова Б.Е., Оспанова А.К., Уваров Н.Ф.</i> Инженерлік технологияда белгілі бір мақсатты қасиетке ие перспективалық композитті материалдарды алу үшін мультижинақтау әдісін (LBL) қолдану (орыс тілінде).....	100
<i>Үмбетова А.К., Слан Г.О., Омарова А.Т., Бурашева Г.Ш., Абидкулова К.Т.</i> Алматы өніріндегі <i>Atrapaxis virgata</i> өсімдігінің химиялық құрамын зерттеу (қазақ тілінде).....	109
<i>Көмекбай Ж.Н., Халменова З.Б., Үмбетова А.К., Бисенбай А.Ф.</i> <i>Melissa officinalis L</i> өсімдігі негізінде фитохимиялық анализ жасау және биологиялық белсенді кешен алу (орыс тілінде).....	114
<i>Надиров Н.К., Некрасов В.Г., Солодова Е.В., Срымов Т., Суханбердиева Д.Т., Култоре М.А.</i> Жаңа буын жылышайы (орыс тілінде).....	122

**СОДЕРЖАНИЕ**

<i>Ергожин Е.Е., Бектенов Н.А., СенГупта Аруп К., Байдулаева А.К., Садыков К.А., Абдрадиева Г.Е., Калмуратова К.М., Рыспаева С.Б.</i> Сорбция ионов стронция новыми комплексообразующими ионитами на основе эпоксиакрилатов и комплексонов (на английском языке).....	6
<i>Аугелханкызы М., Славинская Н.А., Шабанова Т.А., Мансуров З.А.</i> Моделирование окисления и пиролиза аллена (на английском языке).....	12
<i>Ешжанов А.А., Волненко А.А., Левданский А.Э., Корганбаев Б.Н.</i> К расчету эквивалентного диаметра комбинированной регулярно–взвешенной насадки (на английском языке).....	20
<i>Жумадуллаев Д.К., Ешжанов А.А., Волненко А.А., Левданский А.Э.</i> Единый подход к расчету гидравлического сопротивления трубчатого пучка смесительного и поверхностного теплообменников (на английском языке) .....	25
<i>Савденбекова Б.Е., Оспанова А.К., Уваров Н.Ф.</i> Применение метода мультислойной сборки (LBL) в инженерных технологиях для получения перспективных композитных материалов с целенаправленными свойствами (на английском языке).....	31
<i>Альчинбаева О., Сарбаева К.</i> Негативное влияние химических отбеливателей на организм человека (на английском языке).....	38
<i>Умбетова А.К., Слан Г.О., Омарова А.Т., Бурашева Г.Ш., Абидкулова К.Т.</i> Исследование химического состава <i>Atrapaxis virgata</i> алматинского региона (на английском языке).....	42
<i>Баев А.Б., Нурдиллаева Р.Н., Тащенбаева Н.Ж., Озлер М.А.</i> Растворение нержавеющей стали при поляризации переменным током (на английском языке).....	46
<i>Комекбай Ж.Н., Халменова З.Б., Умбетова А.К., Бисенбай А.Г.</i> Фитохимический анализ и разработка получения биологически активного комплекса на основе сырья <i>Melissa officinalis L</i> (на английском языке).....	53
<i>Мамырбекова А., Баев А.Б., Касымова М.К., Мамырбекова А.</i> Влияние перфтордекалина с газотранспортной функцией на рост микроорганизмов (на английском языке).....	59
<i>Еспанова И.Д., Жусупова Л.А., Тапалова А.С., Анназов Н.О.</i> Микроволновая активация реакции присоединения гексен-1 и бутановой кислоты (на английском языке).....	63
<i>Надиров Н.К., Некрасов В.Г., Солодова Е.В., Срымов Т., Суханбердиева Д.Т., Култоре М.А.</i> Теплицы нового поколения (на английском языке).....	70

\* \* \*

<i>Ергожин Е.Е., Бектенов Н.А., СенГупта Аруп К., Байдулаева А.К., Садыков К.А., Абдрадиева Г.Е., Калмуратова К.М., Рыспаева С.Б.</i> Сорбция ионов стронция новыми комплексообразующими ионитами на основе эпоксиакрилатов и комплексонов (на русском языке) .....	81
<i>Ешжанов А.А., Волненко А.А., Левданский А.Э., Корганбаев Б.Н.</i> К расчету эквивалентного диаметра комбинированной регулярно–взвешенной насадки (на русском языке).....	87
<i>Жумадуллаев Д.К., Ешжанов А.А., Волненко А.А., Левданский А.Э.</i> Единый подход к расчету гидравлического сопротивления трубчатого пучка смесительного и поверхностного теплообменников (на русском языке) .....	93
<i>Савденбекова Б.Е., Оспанова А.К., Уваров Н.Ф.</i> Применение метода мультислойной сборки (LBL) в инженерных технологиях для получения перспективных композитных материалов с целенаправленными свойствами (на русском языке).....	100
<i>Умбетова А.К., Слан Г.О., Омарова А.Т., Бурашева Г.Ш., Абидкулова К.Т.</i> Исследование химического состава <i>Atrapaxis virgata</i> алматинского региона (на казахском языке).....	109
<i>Комекбай Ж.Н., Халменова З.Б., Умбетова А.К., Бисенбай А.Г.</i> Фитохимический анализ и разработка получения биологически активного комплекса на основе сырья <i>Melissa officinalis L</i> . (на русском языке).....	114
<i>Надиров Н.К., Некрасов В.Г., Солодова Е.В., Срымов Т., Суханбердиева Д.Т., Култоре М.А.</i> Теплицы нового поколения (на русском языке).....	122

## CONTENTS

<i>Ergozhin E.E., Bektenov N.A., SenGupta Arup K., Baidullaeva A.K., Sadykov K.A., Abdralieva G. E., Kalmuratova K.M., Ryspaeva S.B.</i> Sorption of ions strontium with new complex - forming ionites on the basis of epoxyacrylates and Complexones (in English).....	6
<i>Auyelkhankzy M., Slavinskaya N., Shabanova T.A., Mansurov Z.</i> A modeling study of allene oxidation and pyrolysis (in English).....	12
<i>Yeshzhanov A.A., Volnenko A.A., Levanskii A.E., Korganbayev B.N.</i> To calculating the equivalent diameter of a combined regular-suspended packing (in English).....	20
<i>Zhumadullaev D.K., Yeshzhanov A.A., Volnenko A.A., Levanskii A.E.</i> Common approach to the calculation of hydraulic resistance of a tube bank of contact and surface heat exchangers (in English).....	25
<i>Saydenbekova B.E., Ospanova A.K., Uvarov N.F.</i> Application of the multilayer assembly (LBL) method in engineering technologies for obtaining perspective composite materials with purpose properties (in English).....	31
<i>Alchinbayeva O., Sarbayeva K.</i> Negative effect of chemical bleachers on the human organism (in English).....	38
<i>Umbetova A.K., Slan G.O., Omarova A.T., Burasheva G.Sh., Abidkulova K. T.</i> The study of chemical composition of <i>Atrapaxis virgata</i> from the almaty region (in English) .....	42
<i>Bayeshov A.B., Nurdillayeva R.N., Tashkenbayeva N.Zh., Ozler M.A.</i> Dissolution of stainless steel under alternating current polarization (in English) .....	46
<i>Komekbay Zh. N., Halmenova Z. B., Umbetova A. K., Bisenbay A.G.</i> Phytochemical analysis and development of production of biologically active complex on the basis of raw <i>Melissa officinalis L</i> (in English).....	53
<i>Mamyrbekova A., Bayeshov A.B., Kasymova M.K., Mamyrbekova A.</i> Influence of perfluorodecalin with gas transport function on growth of microorganisms (in English).....	59
<i>Yespanova I.D., Zhusupova L.A., Tapalova A.S., Appazov N.O.</i> Microwave activation of addition of 1-hexene and butanoic acid reaction (in English) .....	63
<i>Nadirov N.K., Nekrasov V.G., Solodova Y.V., Srymov T., Suhanberdieva D.T., Kultore M.A.</i> Hothouses of new generation (in English).....	70

\* \* \*

<i>Ergozhin E.E., Bektenov N.A., SenGupta Arup K., Baidullaeva A.K., Sadykov K.A., Abdralieva G. E., Kalmuratova K.M., Ryspaeva S.B.</i> Sorption of ions strontium with new complex - forming ionites on the basis of epoxyacrylates and complexes (in Russian) .....	81
<i>Yeshzhanov A.A., Volnenko A.A., Levanskii A.E., Korganbayev B.N.</i> To calculating the equivalent diameter of a combined regular-suspended packing (in Russian) .....	87
<i>Zhumadullaev D.K., Yeshzhanov A.A., Volnenko A.A., Levanskii A.E.</i> Common approach to the calculation of hydraulic resistance of a hollow beam of contact and surface heat exchangers (in Russian).....	93
<i>Saydenbekova B.E., Ospanova A.K., Uvarov N.F.</i> Application of the multilayer assembly (LBL) method in engineering technologies for obtaining perspective composite materials with purpose properties (in Russian).....	100
<i>Umbetova A.K., G.O. Slan, Omarova A.T., Burasheva G.Sh., Abidkulova K. T.</i> The study of chemical composition of <i>Atrapaxis virgata</i> from the almaty region (in Kazakh).....	109
<i>Komekbay Zh. N., Halmenova Z. B., Umbetova A. K., Bisenbay A.G.</i> Phytochemical analysis and development of production of biologically active complex on the basis of raw <i>Melissa officinalis L</i> (in Russian).....	114
<i>Nadirov N.K., Nekrasov V.G., Solodova Y.V., Srymov T., Suhanberdieva D.T., Kultore M.A.</i> Hothouses of new generation (in Russian).....	122

**Publication Ethics and Publication Malpractice  
in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct ([http://publicationethics.org/files/u2/New\\_Code.pdf](http://publicationethics.org/files/u2/New_Code.pdf)). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации  
в журнале смотреть на сайте:

www:nauka-nanrk.kz

<http://www.chemistry-technology.kz/index.php/ru/>

**ISSN 2518-1491 (Online), ISSN 2224-5286 (Print)**

Редакторы: *М. С. Ахметова, Т. А. Апендиев, Аленов Д.С.*  
Верстка на компьютере *А.М. Кульгинбаевой*

Подписано в печать 13.02.2018.  
Формат 60x881/8. Бумага офсетная. Печать – ризограф.  
7,8 п.л. Тираж 300. Заказ 1.

---

*Национальная академия наук РК  
050010, Алматы, ул. Шевченко, 28, т. 272-13-18, 272-13-19*